

HOT BARYONS AND THE DISTRIBUTION OF METALS IN THE INTERGALACTIC MEDIUM

JASON TUMLINSON¹ & TAOTAO FANG²*Draft version February 2, 2008*

ABSTRACT

We use the observed number and column-density distributions of intergalactic O VI absorbers to constrain the distribution of metals in the low-redshift intergalactic medium (IGM). In this simple model the metals in the O VI absorbers are assumed to be produced in and propagated from low-redshift galaxies drawn from a real sample, in this case the Sloan Digital Sky Survey (SDSS). This model can explain the observed dN/dz of metals seen in O VI absorbers if these metals are dispersed out to ~ 200 kpc by galaxies down to $L \sim 0.01 - 0.1L_r^*$. Massive galaxies ($L \sim L_r^*$) by themselves cannot provide the necessary enrichment unless they can enrich volumes out to $\gtrsim 0.5 - 1$ Mpc. Alternatively, our model allows an estimate of the fraction of O VI absorbers directly caused by galaxies rather than hot IGM. With this assumption we explore the possible connections between the intergalactic O VI absorbers and the known populations of highly-ionized high-velocity clouds (HVCs) surrounding the Milky Way. Our model predicts that more sensitive, complete surveys optimized to uncover weaker O VI absorbers will find the tentative turnover below $\log N(\text{O VI}) \sim 13.5$ to be a real effect resulting from the apparently limited volumes over which galaxies can enrich the IGM. If so, it would indicate that metals are not as widespread throughout the low-density IGM as they are assumed to be in cosmological simulations of the WHIM.

Subject headings: intergalactic medium — quasars: absorption lines

1. INTRODUCTION

Substantial populations of intergalactic QSO absorption-line systems have been detected in highly-ionized oxygen (O VI) with the Hubble Space Telescope (HST) and the *Far Ultraviolet Spectroscopic Explorer* (*FUSE*). Because they are probably hot ($T \gtrsim 10^5$ K), these absorbers may represent a portion of the “missing baryons” that are predicted (Cen & Ostriker 1999; Davé et al. 1999) to arise at low redshift from continuous structure formation in the warm-hot intergalactic medium (WHIM). Early surveys of these absorbers with HST (Tripp, Savage, & Jenkins 2000; Savage et al. 2002, Tripp 2005) found unexpectedly high number densities per unit redshift, $dN/dz \sim 16$, down to 30 - 50 mÅ sensitivity. The recent *FUSE* study by Danforth & Shull (2005, hereafter DS05) has refined these estimates of dN/dz and measured their column density distribution, $dN/dNdz$. DS05 assumed uniform 0.1 solar metallicity ($[\text{O}/\text{H}] = -1.0$) and $T = 10^{5.5}$ K to estimate the total contribution of the O VI absorbers to the baryon density, $\Omega_{\text{OVI}}h_{70}^{-1} = 0.0022 \pm 0.0003$, where $\Omega_b h_{70}^2 = 0.024 \pm 0.001$ (Spergel et al. 2003)³.

The *FUSE* data cannot measure the gas metallicity precisely but they allow for estimates of typical values. DS05 detect 40 O VI absorbers which span 2.5 dex in $N(\text{H I})$ but only 1.5 dex in $N(\text{O VI})$. By combining photoionization and collisional ionization equilibrium (CIE) models with a theoretical relation between IGM density and N_{HI} , DS05 derive a typical metallicity of $Z_{\text{O}} = 0.09Z_{\odot}$. We derive a conservative lower limit $[\text{O}/\text{H}] = -2.3 \pm 0.05$ from a pure CIE model that matches

the broad locus of points in Figure 1 of DS05, assuming the maximum fractional ionization, $f_{\text{OVI}} = 0.2$, of O VI at $T = 10^{5.45}$ K. However, according to Equation 6 of DS05 this metallicity limit implies $\Omega_{\text{OVI}}h_{70}^{-1} = 0.044$, close to the total cosmic baryon density. Because a major portion of the WHIM is thought to lie in hotter gas accessible only to X-ray telescopes (Fang et al. 2002; Mathur, Weinberg, & Chen 2003; McKernan et al. 2003; Nicastro et al. 2005), it is likely that the true metallicity of these absorbers lies closer to the fiducial value $[\text{O}/\text{H}] = -1.0$ assumed by DS05. Metallicity $[\text{O}/\text{H}] \gtrsim -1.0$ in $\sim 30\%$ of the detected Ly α clouds (DS05) is too high and too common to have been in place by $z \sim 3$, where Schaye et al. (2003) find filling factors of 2 - 10% for gas with $[\text{C}/\text{H}] \geq -2.0$. The O VI absorbers therefore imply significant later enrichment by galaxies. Conversely, if the metallicities of these absorbers lie near our lower limits, a portion of the sample would need to be removed to limit their cosmic density. Thus the mean metallicity, the dispersion in the metallicity, and the possibility of other mechanisms for producing O VI are essential to understanding the contribution of O VI absorbers to Ω_b .

To produce detectable O VI absorption, an intervening cloud must (1) have the proper ionization conditions to ionize oxygen to O VI, and (2) have sufficient metallicity to make the O VI lines strong enough to detect. The first condition can be met in CIE gas produced in shocks (Shull & McKee 1979) or in hot-cold interfaces (Fox et al. 2004; Indebetouw & Shull 2004, and references therein), though for some IGM conditions photoionization by a metagalactic ionizing background (Haardt & Madau 1996) can also produce O VI. The ionization mechanism need not be caused by, nor occur near, a galaxy, but in the standard picture of structure formation metals must be created in galaxies. When absorber/galaxy relationships are considered, there is no necessary connection between galaxies and the *ionization*

¹ Department of Astronomy and Astrophysics, University of Chicago, 5640 S. Ellis Ave., Chicago, IL 60637

² Chandra Fellow, Astronomy Department, 601 Campbell Hall, University of California, Berkeley, CA 94720

³ We assume a Hubble constant $H_0 = (70 \text{ km s}^{-1} \text{ Mpc}^{-1}) h_{70}$.

of the gas, while there is a necessary connection between galaxies and the *metallicity* of the gas. Thus even if the O VI absorbers fill only a portion of the IGM hot-gas budget, they can still constrain the distribution of metals. In this paper we study the dispersal of IGM metals by galaxies using only *observables* and simple geometric arguments. Our model does not attempt to explain the shock-heating or photoionization mechanisms that cause or maintain the ionization of O VI, but rather the metals that must have some link to nearby galaxies.

Our model has significant advantages over simulations (Cen et al. 2002; Fang & Bryan 2001; Chen et al. 2003; Yoshida et al. 2005) or semi-analytic calculations (Furlanetto et al. 2005) that predict dN/dz from *a priori* models. These models either do not treat metal distribution in a self-consistent way, by adopting a uniform or density-dependent metallicity, or they use complicated models to calculate ionization structures and obtain reasonable O VI ionization fractions. By contrast, our model avoids all these complicated issues. By decoupling these entangled physical processes, we can then focus on key issues, namely the volume that can be enriched by galaxies, and the minimum luminosity of the galaxies that can enrich the nearby IGM. Our method also allows us to evaluate the hypothesis that the intergalactic O VI absorbers are related to the O VI HVCs seen in the Galaxy. Combining some simple geometrical arguments we derive the expected column density distribution, $dN/dNdz$. When compared with observations, our simple model provides significant constraints on the distribution of metals and their connection with galaxies.

2. QSO/GALAXY PAIRS AND IGM METAL ENRICHMENT

Our model compares the distribution of IGM metals expressed by dN/dz to the incidence of QSO-galaxy “pairs” expected per redshift interval. We calculate the total number of absorbers dn that can be encountered by one random sightline along a distance dl to a background QSO. The spatial density of halos with absolute magnitude between M_r and $M_r + dM_r$ is $(dn_h/dM_r)dM_r$, where dn_h/dM_r will ultimately be determined by SDSS. Assuming each halo can enrich the surrounding volume to a radius of R , the total number of such intercepted absorbers within a distance of dl along the line of sight (LOS) toward background quasars then is

$$\frac{dn}{dl} = f\pi R^2 \int_{-\infty}^{M_r^{faint}} \frac{dn}{dM_r} dM_r. \quad (1)$$

Here f is the LOS covering fraction, representing the chances of intercepting absorbers within each halo. In the next section we will see that $f \sim 1$ when we relate the local O VI high velocity clouds (HVCs) and the intergalactic O VI absorbers.

In principle the integral in the last term of Equation 1 could be stated in terms of mass and derived from, e.g. a Press-Schechter function, but in practice we use observations. We begin with the SDSS Data Release 3 catalog of QSOs (all spectroscopically confirmed) and galaxies⁴. To maximize the size of the sample and therefore provide good statistics, we use all SDSS photometric sources determined to be galaxies by photometric selection, then calculate photometric redshifts, z_{photo} , from

the SDSS *ugriz* photometry using the software of Blanton et al. (2003). These redshifts have typical errors $\sigma_z = 0.02 - 0.04$, which make individual pairs quite uncertain but which wash out in the average. For each QSO sightline, we assign to all galaxies within $42'$ an impact parameter b to the QSO sightline at the galaxy z_{photo} . We require $z_{\text{photo}} = 0.02 - 0.2$ to minimize spurious detections on the low end and incompleteness at the high end. We correct for incompleteness in faint galaxies at the higher redshifts by assigning each galaxy (or magnitude bin of galaxies) a unique pathlength Δz based on its absolute magnitude and the SDSS flux limits. Then, for a given spherical “halo” or “enrichment” radius R and a LOS covering fraction f , we can derive the total number and dN/dz of pairs, with cuts for any galaxy property (i.e., color, morphology, luminosity) measurable from the *ugriz* photometry and z_{photo} .

Figure 1 shows differential and cumulative distributions of dN/dz achieved by galaxies of varying L_r , given a uniform impact parameter b and $f = 1$. The key measurements are $dN/dz = 3, 9, 17, 19$, and 21 for $W_\lambda \geq 100$, $50, 30, 15$, and 10 mÅ, respectively (DS05). This mapping between galaxy populations and absorber incidence is the major result of this study. Spherical enrichment zones must be quite large, $b \gtrsim 0.5$ Mpc, and uniform ($f = 1$) for enrichment by massive galaxies ($L \gtrsim L_r^*$) to explain the observed numbers of O VI absorbers. For $b \sim 200$ kpc, low-mass galaxies ($L \sim 0.1L_r^*$) can reproduce the observed dN/dz . To obtain $dN/dz = 21$ with dwarf galaxies only, we require uniform $R = 145$ kpc for $M_r = -19$ to -16 and $R = 160$ kpc for $M_r = -18$ to -16 . In passing we note our model can also constrain R for H I halos corresponding to high- $N(\text{H I})$ Ly α clouds, i.e. $R = 50$ kpc at $L \gtrsim L_r^*$ is needed to reproduce $dN/dz = 1$ for Lyman limit systems (Keeney et al. 2005).

DS05 find a power-law dependence of dN/dz on N_{OVI} , $dN/dN_{\text{OVI}} \propto N_{\text{OVI}}^{-2.2}$ for absorbers with $\log N(\text{O VI}) \geq 13.4$. The behavior of $dN/dNdz$ at low N is of critical importance to estimating Ω_{OVI} . However, below $\log N(\text{O VI}) = 13.4$, the observed numbers of O VI absorbers drop below the power-law fit, and well below the numbers expected from simulations that assume uniform or overdensity-dependent metallicity in the WHIM (Figure 2). DS05 cannot say whether the apparent turnover below $\log N(\text{O VI}) \simeq 13.5$ is real or due to survey incompleteness below ~ 50 mÅ. Integrating this power law down to $\log N(\text{O VI}) = 13.0$ (12.5 mÅ) yields $dN/dz = 106$. This high number density requires $[\text{O}/\text{H}] \gtrsim -2$ over ~ 600 kpc spheres for $L \gtrsim 0.1L_r^*$ (Figure 1), an unlikely outcome. We believe it is more likely that the observed turnover below $\log N(\text{O VI}) \simeq 13.5$ is real and not a consequence of incompleteness in the DS05 survey. This prediction could be verified by a survey with more pathlength coverage down to $\log N(\text{O VI}) = 13.0$ (12.5 mÅ). If this deficit relative to theoretical expectations is real, our model suggests that metals are not as widely or uniformly distributed throughout the low-density IGM as simulations assume or find them to be.

QSO absorption-line studies of the dispersal of IGM metals generally offer conflicting guidance because they suffer from the practical difficulty of identifying the faintest galaxy with which a given absorber might be associated. These studies cannot generally exclude the

⁴ Available at <http://skyserver.pha.jhu.edu>.




FIG. 1.— Upper panel: Differential distribution of dN/dz as a function of galaxy luminosity, in 1 mag bins. From top, the lines correspond to $R = 500, 300$, and 100 kpc. Lower panel: As a function of b and SDSS r -band absolute magnitude, contours of constant cumulative dN/dz obtained in the fiducial model ($f = 1$). Clearly dwarf galaxies must enrich their environments for the observed number density of metal-line absorbers to be obtained. The vertical lines mark L^* and $0.1L^*$ from the Blanton et al. (2003) SDSS luminosity functions. Solid contours for $dN/dz = 3, 9, 17$, and 21 correspond to equivalent width limits $W_\lambda = 100, 50, 30$, and 10 mÅ, respectively (DS05). The dashed contours show model predictions for higher dN/dz . The error bar at the lower left shows the approximate vertical range associated with the errors bars in the observed dN/dz . The symbols mark the luminosities and impact parameters of nearest-neighbor galaxies for seven well-studied QSO absorbers. These are: open circle, H1821+643 (Tripp et al. 2001); filled circles, PG0953+415 (Savage et al. 2002); open triangle, PKS2155-304 (Shull, Tumlinson, & Giroux 2003); diamond, PKS0405-419 (Prochaska et al. 2005); open squares, PG1211+143 (Tumlinson et al. 2005).

presence of a faint galaxy nearer the sightline than a known bright one, and so they tend to overestimate the dispersal of metals. This effect is seen in the data points in Figure 1, which show M_r and b of nearest-neighbor galaxies to O VI absorbers with galaxy surveys of varying depth and sky coverage. There is a general trend of metals spread to $\sim 100 - 200$ kpc, with some notable exceptions. The $z = 0.06807$ absorber toward PG0953+415 (Savage et al. 2002) suggests $\gtrsim 0.2$ solar enrichment to ~ 600 kpc, but this field has not been surveyed much below L^* . The PKS2155-304 O VI absorbers (Shull, Tumlinson, & Giroux 2003) provide the best case for metal dispersal to ~ 400 kpc, because this field has been surveyed to $L \sim 0.01L^*$, but the absorbers appear to be associated with a galaxy group that may complicate constraints on metal dispersal. Chen et al. (2001) systematically studied C IV near 50 galaxies over a range of luminosity and found $f = 0.67$ at $b \leq 100$ kpc and $f = 0.06$ at $b \geq 100$ kpc, with typical limiting $W_\lambda \sim 100$ mÅ. This

result does not rely on a post-facto galaxy survey, and so provides more robust evidence that luminous galaxies have associated metals out to $b \sim 100$ kpc. Stocke et al. (2004) found a post-starburst dwarf with $M_B = -13.9$ associated with a 6% solar metallicity absorber in the sightline to 3C 273. This galaxy, while starbursting, apparently drove a metal-enriched superwind $71h_{70}^{-1}$ kpc into the IGM. Thus there is evidence that metal enrichment spreads to ~ 100 kpc from all galaxies, but evidence for further propagation is less clear.

We stress that the known dN/dz of O VI absorbers represents the minimum budget of metals that must be explained in terms of enrichment by galaxies. There are surely other reservoirs of metals that meet condition 2 above (detectable metallicity) but not 1 (conditions that produce O VI). Any difficulty associated with distributing the oxygen seen in O VI can only be compounded by the addition of these other reservoirs (e.g., photoionized gas in C IV systems) to the budget of IGM metals. Finally, Figure 1 shows results for $f = 1$ and so presents the most optimistic case of enrichment.

3. INTERGALACTIC O VI AND HIGH-VELOCITY CLOUDS

Our model can also interpret the intergalactic O VI absorbers as analogs of the local highly-ionized high-velocity clouds (Sembach et al. 2003, hereafter S03). To do so, we must first derive a mapping between the covering fraction, C , of HVCs with column density N as seen along radial sightlines out from the Galaxy, to f , the covering fraction along a LOS that follows a chord through an intervening galaxy halo. The key assumption here is that the HVCs are not *unique*, i.e., an observer located in another galaxy similar to our Galaxy should be able to detect similar HVCs with a similar pattern of column density distribution. Let us first consider HVCs with column densities between N_j and $N_j + \Delta N_j$, and with a covering fraction C_j – the total covering fraction C is the sum of C_j . Assuming there are a total of T_j such absorbers uniformly distributed within the halo (of a galaxy or a group), and each absorber has a spherical geometry with radius d , we find T_j and the spatial number density of the absorbers within the halo n_j are

$$T_j = \frac{4R^2}{3d^2} C_j; \quad n_j \equiv \frac{T_j}{V} = \frac{C_j}{\pi d^2 R}, \quad (2)$$

respectively, where R is the radius of the halo. These absorbers project onto the sky a total area $A_j = T_j \pi d^2$, such that their covering fraction along the LOS is

$$f_j = \frac{A_j}{\pi R^2} = \frac{4}{3} C_j. \quad (3)$$

We derive all quantities from observation – C_j from Table 4 of S03 and the spatial density of galaxies from SDSS. The total LOS covering fraction f then is $f = \sum_j f_j = (4/3)C$. Adopting the value of C from S03, we find f ranges from ~ 0.64 to ~ 1.1 down to 30 mÅ, which justifies the value of 1 we adopt in the previous section. Our results appear in Figure 2, where we show that HVC-like O VI absorbers can make a significant contribution to the observed numbers if massive galaxies have large halos, or if all galaxies down to $M_r \sim -16$ ($\sim 0.01L^*$) have O VI HVCs out to ~ 100 kpc. Highly ionized gas can arise in galaxy superwinds (Heckman et al. 2001), in gas tidally stripped from galaxies (Cox et




FIG. 2.— Cumulative dN/dz of O VI absorbers as a function of W_λ for the DS05 data points (filled diamonds) and theory. The solid line marks the prediction for the WHIM simulation of Fang & Bryan (2001), with an overdensity-dependent metallicity. The dotted and dashed lines mark the present model for dN/dz with column-density dependent covering fractions $f_j = 4C_j/3$ from S03, for luminosity-impact parameter pairs $(-19, 100)$ and $(-16, 80)$, respectively. Note that HVC-type O VI absorbers may make significant contributions to the observed numbers, especially if they arise from dwarf galaxies.

al. 2004), and in accreting gas (Maller & Bullock 2004; Birnboim & Dekel 2002). The $M_B = -13.9$ dwarf $71h_{70}^{-1}$ kpc away from a metal-enriched absorber toward 3C 273 (Stocke et al. 2004) is one example at the faint end of the luminosity function. The *FUSE* studies of Galactic O VI HVCs (Sembach et al. 2003; Fox et al. 2004; Collins, Shull, Giroux 2004) show that O VI can arise in the extended halo regions of larger galaxies (~ 100 kpc) and not necessarily correspond to the WHIM. From Figure 2, it appears that if these processes have high f out to ~ 100 kpc around galaxies down to $\sim 0.01L_r^*$, they may contribute $dN/dz \sim 5$ to the overall population of O VI absorbers. Given the potentially important contributions of O VI absorbers to Ω_{WHIM} , their utility as pointers to the hotter WHIM seen in X-ray absorption, and considering the many mechanisms that can produce O VI near galaxies, it is important to obtain an unbiased observation of the incidence of O VI and metals within 100–200 kpc of galaxies over a range of mass. Until such an observation is made, we cannot know how to correct the inferred baryon densities for these contaminating effects. Our observations can also guide the design of such

studies by constraining the fields over which galaxy surveys must be executed.

4. CONCLUSIONS

Based on our model of the distribution of IGM metals and the observed numbers of IGM O VI absorbers, we draw the following conclusions:

1. Unless they eject metals to $\sim 0.5 - 1$ Mpc, luminous galaxies cannot explain dN/dz for metal-line absorbers in the low- z IGM. If low-mass galaxies ($\gtrsim 0.01 - 0.1L_r^*$) disperse metals to ~ 200 kpc, dN/dz for the O VI absorbers can be explained.
2. Uniform enrichment to ~ 1 Mpc is implied by integrating the power-law dependence of $dN/dNdz$ (DS05) to $\log N(\text{O VI}) = 13.0$. Thus the turnover seen by DS05 in $dN/dNdz$ below $\log N(\text{O VI}) \simeq 13.5$ (50 mÅ) is more likely a real effect than a consequence of survey incompleteness. A survey sensitive to $W_\lambda = 12.5$ mÅ could confirm this result and further constrain the dispersal of IGM metals. Conversely, additional absorbers below $\log N(\text{O VI}) \simeq 13.5$ would imply that metals are quite widespread.
3. A high covering fraction of O VI from “HVCs” within 100–200 kpc from galaxies can contribute $dN/dz = 5 - 10$, a non-negligible portion of the observed total that would not contribute to Ω_{WHIM} .

Thanks to Charles Danforth and Mike Shull for sharing their results before publication and to Charles Danforth and John Stocke for constructive comments. We began this work in the hospitality of the Kavli Institute for Theoretical Physics at the University of California, Santa Barbara, supported in part by NSF under Grant No. PHY99-07949. J. T. is supported by the Astronomy and Astrophysics Department at the University of Chicago and HST GO-9874.01-A. T. F. was supported by NASA through *Chandra* Postdoctoral Fellowship PF3-40030 issued by the *Chandra* X-ray Observatory Center, which is operated by the Smithsonian Astrophysical Observatory for and on behalf of NASA under contract NAS8-39073.

REFERENCES

- Birnboim, Y., & Dekel, A. 2002, MNRAS, 345, 349
 Blanton, M. R., et al. 2003, ApJ, 592, 819
 Cen, R., & Ostriker, J. P. 1999, ApJ, 519, L109
 Chen, X., Weinberg, D. H., Katz, N., Davé, R. 2003, ApJ, 594, 42
 Collins, J. A., Shull, J. M., & Giroux, M. L. 2004, ApJ, 605, 216
 Cox, T. J., et al. 2004, ApJ, 607, L87
 Danforth, C. W., & Shull, J. M. 2005, ApJ, in press
 Fang, T., & Bryan, G. L. 2001, ApJ, 561, L31
 Fang, T., et al. 2002, ApJ, 572, 127
 Fox, A. J., et al. 2004, ApJ, 602, 738
 Furlanetto, S., Phillips, L. A., & Kamionkowski, M. 2005, MNRAS, in press
 Haardt, F., & Madau, P. 1996, ApJ, 461, 20
 Heckman, T., et al. 2001, ApJ, 554, 1021
 Indebetouw, R., & Shull, J. M. 2004, ApJ, 605, 205
 Keeney, B. A., et al. 2005, ApJ, submitted (astro-ph/0409448)
 Maller, A. H., & Bullock, J. S. 2004, MNRAS, 355, 694
 Mathur, S., Weinberg, D. H., & Chen, X. 2003, ApJ, 582, 82
 McKernan, B., et al. 2003, ApJ, 583, L83
 Nicastro, F., et al. 2005, Nature, in press
 Prochaska, J. X., Chen, H.-W., Howk, J. C., Weiner, B. J., & Mulchaey, J. 2004, ApJ, 617, 718
 Savage, B. D., et al. 2002, ApJ, 564, 631
 Schaye, J., Aguirre, A., Kim, T.-S., Theuns, T., Rauch, M., & Sargent, W. L. W. 2003, ApJ, 596, 768
 Shull, J. M., & McKee, C. F. 1979, ApJ, 227, 131
 Shull, J. M., Tumlinson, J., & Giroux, M. L. 2003, ApJ, 594, L107
 Spergel, D. N., et al. 2003, ApJ, 148, 175
 Stocke, J. T., et al. 2004, ApJ, 609, 94
 Tripp, T. M., Savage, B. D., & Jenkins, E. B. 2000, ApJ, 534, L1
 Tripp, T. M., Giroux, M. L., Stocke, J. T., Tumlinson, J., & Oegerle, W. R. 2001, ApJ, 563, 724
 Tripp, T. M. 2005, in Astrophysics in the Far Ultraviolet, in press
 Tumlinson, J., Shull, J. M., Giroux, M. L., & Stocke, J. T. 2005, ApJ, 620, in press
 Yoshida, N., Furlanetto, S. R., & Hernquist, L. 2005, ApJ, 618, 91